

A Variant in *LIN28B* Is Associated with 2D:4D Finger-Length Ratio, a Putative Retrospective Biomarker of Prenatal Testosterone Exposure

Sarah E. Medland,^{1,9} Tetyana Zayats,^{2,3,9} Beate Glaser,^{2,3} Dale R. Nyholt,¹ Scott D. Gordon,¹ Margaret J. Wright,¹ Grant W. Montgomery,¹ Megan J. Campbell,¹ Anjali K. Henders,¹ Nicholas J. Timpson,^{2,3} Leena Peltonen,^{4,5,6,7} Dieter Wolke,⁸ Susan M. Ring,³ Panos Deloukas,⁴ Nicholas G. Martin,¹ George Davey Smith,^{2,3} and David M. Evans^{2,3,*}

The ratio of the lengths of an individual's second to fourth digit (2D:4D) is commonly used as a noninvasive retrospective biomarker for prenatal androgen exposure. In order to identify the genetic determinants of 2D:4D, we applied a genome-wide association approach to 1507 11-year-old children from the Avon Longitudinal Study of Parents and Children (ALSPAC) in whom 2D:4D ratio had been measured, as well as a sample of 1382 12- to 16-year-olds from the Brisbane Adolescent Twin Study. A meta-analysis of the two scans identified a single variant in the *LIN28B* gene that was strongly associated with 2D:4D (rs314277: $p = 4.1 \times 10^{-8}$) and was subsequently independently replicated in an additional 3659 children from the ALSPAC cohort ($p = 1.53 \times 10^{-6}$). The minor allele of the rs314277 variant has previously been linked to increased height and delayed age at menarche, but in our study it was associated with increased 2D:4D in the direction opposite to that of previous reports on the correlation between 2D:4D and age at menarche. Our findings call into question the validity of 2D:4D as a simplistic retrospective biomarker for prenatal testosterone exposure.

Introduction

The ratio of the lengths of the second to fourth digits (2D:4D) is a sexually dimorphic trait that is on average a quarter of a standard deviation lower in males than in females.¹ First identified by Ecker in 1875;² the measure was rediscovered by Wilson in the early 1980s³ and subsequently by Manning, who hypothesized that the ratio reflected prenatal androgen exposure.^{4,5} Consistent with this theory, sex differences in 2D:4D develop prenatally⁶ and remain relatively stable across the lifespan.⁷ Given the practical and ethical difficulties inherent in measuring testosterone exposure in the developing fetus, many researchers have adopted 2D:4D as a noninvasive retrospective biomarker for prenatal androgen exposure, although its use as such is controversial (see McIntyre et al. for a review⁸). Despite this controversy, over 300 papers using this measure have been published in the last 10 years,⁹ and 2D:4D has been shown to correlate with a wide range of diseases and physiological and psychological traits, including autism,¹⁰ attention deficit disorder,¹¹ fertility,¹² myocardial infarction,¹³ visuo-spatial ability,¹⁴ homosexuality,^{15,16} athletic performance,¹⁷ and age at menarche.¹⁸

2D:4D is highly heritable, with additive genetic effects explaining ~60% of the phenotypic variance.^{19–22} Although the results from one twin study suggested that female twins exhibit higher heritabilities than males for left hand 2D:4D,¹⁹ a larger study failed to find any significant sex

limitation or differences in the magnitude of heritability between the sexes for 2D:4D traits.²⁰ There has also been little progress identifying the individual variants underlying this genetic variation. Following the hypothesis that 2D:4D reflects prenatal exposure to testosterone, Manning et al.²³ examined the association between the number of CAG repeats at the androgen receptor locus (*AR* [MIM 313700]) and 2D:4D in males ($n = 51$) and reported a significant correlation between CAGn and 2D:4D for the right ($r = 0.29$) but not the left ($r = 0.005$) hand. To the best of our knowledge, no other genetic association studies using this trait have been performed to date.

In order to identify genetic determinants underlying variation in 2D:4D, we applied a genome-wide association approach to study 1507 children from the Avon Longitudinal Study of Parents and Children (ALSPAC), a large population-based cohort in which 2D:4D had been measured at ~11 years of age,²⁴ as well as a sample of 1382 12-, 14-, and 16-year-old twins and their singleton siblings from the Queensland Institute of Medical Research, Brisbane Adolescent Twin Study (QIMR, Australia).^{20,25}

Subjects and Methods

Participants

ALSPAC is a population-based birth cohort study consisting of over 13,000 women and their children recruited

¹Genetic Epidemiology, Queensland Institute of Medical Research, Australia, 4006; ²MRC Centre for Causal Analyses in Translational Epidemiology, University of Bristol BS82BN, UK; ³Department of Social Medicine, University of Bristol, Bristol BS82PS, UK; ⁴Wellcome Trust Sanger, Institute, Cambridge CB10 1SA, UK; ⁵Biomedicum Helsinki, Research Program in Molecular Medicine, University of Helsinki FI-00290, Finland; ⁶Department of Molecular Medicine, National Public Health Institute, Helsinki FI-00271, Finland; ⁷The Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA; ⁸Department of Psychology and Health Sciences, University of Warwick, Coventry CV47AL, UK

⁹These authors contributed equally to this work

*Correspondence: dave.evans@bristol.ac.uk

DOI 10.1016/j.ajhg.2010.02.017. ©2010 by The American Society of Human Genetics. All rights reserved.

from the county of Avon, UK, in the early 1990s.²⁴ Both mothers and children have been extensively followed from the eighth gestational week onward with a combination of self-reported questionnaires, medical records, and physical examinations. Biological samples including DNA have been collected for 10,121 of the children from this cohort. The discovery sample reported in this study concerns 1507 children who had their 2D:4D measured at 11 years of age (mean = 11.75 years) and for whom genome-wide SNP typing had been performed.²⁶ The replication sample consisted of an additional 3659 children from ALSPAC who were not part of the initial discovery cohort.

Participants in the QIMR Brisbane Adolescent Twin Study were recruited from the general population, in the context of ongoing studies of melanoma risk factors and studies of cognition.²⁷ Twins and their singleton siblings were enlisted by contacting the principals of primary schools in the greater Brisbane area, by media appeals, and by word of mouth. It is estimated that approximately 50% of the eligible birth cohort were recruited into the study, which began in 1992. Digit ratios were available for 1382 individuals with genome-wide association data from 671 families (comprising 169 singletons, 332 sibling pairs, 135 sibling trios, 31 sibling quads, and 4 sibling quins). Age range for the sample was 11–24 years (mean = 15.46, standard deviation [SD] = 3.27).

In both samples, participants' hands were photocopied during a clinical visit, and measurements of the second and fourth fingers were taken from the photocopies with the use of digital calipers (accurate to 0.1 mm). The 2D:4D was calculated as the length of the second digit divided by the length of the fourth digit, multiplied by 100 so as to avoid computational difficulties due to the low variance of the trait. In both samples, the measure was normally distributed, so no further transformation was required.

In ALSPAC, a random sample of 57 right and 48 left hands were measured *in vivo* to establish the validity of using a photocopy measurement to assess 2D:4D. Similarly, in the QIMR cohort, 680 hands were measured twice from the same hand photocopy, once by hand with the use of digital calipers and once with the use of a computer-assisted measurement program, and the reliability between measurement occasions was calculated.

Children's standing height in ALSPAC was measured with a Harpenden Stadiometer. Both studies were performed with the approval of the appropriate ethics committees, and informed consent was obtained from all participants and their parents.

Genotyping

A total of 1543 ALSPAC children were initially genotyped at 317,504 SNPs on the Illumina HumanHap 317K SNP chip. Individuals exhibiting cryptic relatedness, non-European ancestry, high genome-wide heterozygosity, and/or missing rates were removed from analyses as described

previously,²⁶ leaving 1507 individuals in the analysis who had been measured for 2D:4D. Markers with minor allele frequency (MAF) < 1%, SNPs with > 5% missing genotypes, and any marker that failed an exact test of Hardy-Weinberg equilibrium (HWE; $p < 5 \times 10^{-7}$) were excluded from further analysis, leaving 310,613 SNPs that passed quality control.

The QIMR participants analyzed here were genotyped on the Illumina Human610-Quad SNP chip. These samples were genotyped in the context of a larger GWAS, which resulted in the genotyping of 16,140 individuals²⁷ with the use of the Illumina 317, 370, and 610 SNP chips. Genotype data were screened for genotyping quality (GenCall < 0.7), SNP and individual call rates (< 0.95), HWE failure ($p < 10^{-6}$), and MAF (< 0.01). Because these samples were genotyped in the context of a larger project, the data were integrated with the larger QIMR genotype project and checked for pedigree, sex and Mendelian errors and for non-European ancestry. Because the QIMR genotyping project included data from the 317, 370, and 610 chip sets, to avoid introducing bias to the imputed data, a set of SNPs common to the three genotyping platforms was used for imputation ($n = 274,604$).

Follow-up genotyping of two SNPs in the *LIN28B* gene (MIM 611044), rs314277 and rs314276, was carried out in an additional 5129 individuals from the ALSPAC cohort by K-Biosciences, who employ a novel form of competitive allele-specific PCR (KASPar) and TaqmanTM system for genotyping. The rs314277 SNP was chosen for replication because it showed maximum association in the discovery cohort, whereas rs314276 was chosen because it had previously shown association with traits correlated with 2D:4D, such as age of menarche.²⁸

Statistical Analyses

Because ALSPAC and QIMR samples were genotyped on different arrays, consensus autosomal genotypic data were imputed with Markov Chain Haplotyping software (MACH) with the use of phased data from CEU individuals from release 22 of the HapMap project as the reference set of haplotypes. Only SNPs that could be imputed with relatively high confidence ($R^2 > 0.3$) and had a MAF > 1% were used in subsequent analyses. In the ALSPAC cohort, association analysis of imputed SNPs was performed assuming an underlying additive model with the use of the software package MACH2QTL, which accounts for uncertainty in prediction of the imputed data by weighting genotypes by their estimated posterior probabilities. In the QIMR twins study, the most likely genotypes were imputed at each locus, and these genotypes were subsequently analyzed with MERLIN.²⁹ Markers at physically genotyped loci on the X chromosome were analyzed with PLINK in the ALSPAC sample³⁰ and MINX in the QIMR cohort.²⁹ SNPs were tested for association with right 2D:4D, left 2D:4D, and the mean of left and right values. All analyses included sex as a covariate. Results for the two cohorts were then combined with the use of fixed-effects inverse

Table 1. Mean and Standard Deviation for 2D:4D in the Discovery and Replication Cohorts

2D:4D Ratio		Mean (SD) 2D:4D in Original GWAS Sample		Mean (SD) 2D:4D in Replication ALSPAC Cohort
		QIMR	ALSPAC	
Right	Females	97.93 (3.18)	96.67 (3.25)	96.73 (3.17)
	Males	96.16 (3.30)	95.85 (3.19)	95.73 (3.19)
Left	Females	98.43 (3.46)	96.80 (3.34)	96.88 (3.16)
	Males	96.84 (3.43)	96.02 (3.23)	95.93 (3.15)
Average	Females	98.18 (2.91)	96.74 (3.04)	96.82 (2.90)
	Males	96.84 (3.44)	95.94 (2.93)	95.84 (2.85)

Females tend to have higher ratios than males.

variance meta-analysis. We also performed a chi-square test for heterogeneity to test whether the regression coefficients differed significantly between males and females for the regression of 2D:4D for all SNPs in the discovery GWAS (i.e., a test for additive genotype \times sex interaction). The p values for the tests of heterogeneity were combined across ALSPAC and QIMR cohorts with the software package METAL to produce an overall level of significance. Association in the replication sample was performed in PLINK via linear regression assuming an underlying additive genetic model.

Results

The Pearson's product moment correlation between the in vivo and photocopied measurements of the length of the second and fourth digits was high for the right and left hands (all $r > 0.97$). The correlations between finger lengths measured with the use of digital calipers and those measured with the use of a computer-assisted program were similarly high (all $r > 0.96$), suggesting that our measurements have a high degree of repeatability.

Table 1 displays the mean and SD of the 2D:4D measurements for the left hand, right hand, and mean of both hands in the ALSPAC and QIMR discovery cohorts, as well as in the ALSPAC replication sample. As expected, across samples and hands, the mean 2D:4D was higher for females than for males. Quantile-quantile plots (Q-Q plots) for the genome-wide association scan of ALSPAC individuals, the GWAS of the QIMR twins, and the combined meta-analyses are presented in Figure S1, available online. Both plots indicate that the observed GWAS test statistics lie close to expectation and suggest that potential technical and stratification artifacts had negligible impact on the results. Consistent with this interpretation, the genomic inflation factors in the ALSPAC ($\lambda = 1.01$), QIMR ($\lambda = 1.01$), and meta-analyzed ($\lambda = 1.00$) samples indicate little inflation of the association test statistics. We also checked whether p values

derived from markers on the X chromosome might exhibit a more dramatic deviation from the null hypothesis of no association than the genome as a whole. Q-Q plots of the combined meta-analyses for left-hand 2D:4D, right-hand 2D:4D, and mean 2D:4D showed that there was little evidence that this was the case, although a few markers on the X chromosome did exceed null expectations for left-hand 2D:4D (Figure S2).

The genome-wide association results for the combined meta-analysis of left-hand, right-hand, and mean 2D:4D are presented in Figures S3–S5. A single SNP, rs314277, in the *LIN28B* gene (Figure 1) reached genome-wide significance for mean 2D:4D ($p = 4.1 \times 10^{-8}$), as well as suggestive significance for left-hand ($p = 1.5 \times 10^{-6}$) and right-hand (8.2×10^{-7}) 2D:4D. Each copy of the minor allele was associated with a 0.6 increase in mean 2D:4D. No other imputed or genotyped SNP in this region met the criterion for genome-wide significance, although several SNPs, including rs314276, which had shown association with pubertal development and height in previous studies,²⁸ showed nominal evidence of association with 2D:4D (left: $p = 7.9 \times 10^{-5}$; right: $p = 1 \times 10^{-3}$; mean: $p = 5.4 \times 10^{-5}$). Interestingly, although the recombination rate in genomic regions 50 kb either side of rs314277 was low, there were few SNPs in appreciable r^2 with the marker, which may explain why the next most associated SNP had p values at least two orders of magnitude lower. Conditioning on rs314277 in both the ALSPAC and QIMR data sets reduced the signal at the surrounding loci but did not completely abolish all evidence of association (ALSPAC best $p = 0.0021$; QIMR best $p = 0.0069$), suggesting either the existence of a more strongly associated variant in the region that had not been imputed or a smaller second signal independent of rs314277 associated with 2D:4D in these data.

There was close correspondence in the topmost hits between left and right 2D:4D, which is not surprising, given the moderate to high phenotypic correlation between these variables (ALSPAC: $r = 0.69$; QIMR $r = 0.56$). We therefore present only the results for the mean 2D:4D in the main text. Tables S1–S3 list all SNPs with a combined p value of less than 1×10^{-5} for the meta-analysis of left-hand, right-hand, and mean 2D:4D. These included SNPs in the genes *SMOC1* (MIM 608488), *SOX7* (MIM 612202), *GLIS1* (MIM 610378), *EFNA1* (MIM 191164), *VAV3* (MIM 605541), and *NEDD4L* (MIM 606384). Of particular note were SNPs on chromosome 8 in the *SOX7* gene, which were strongly associated with 2D:4D in the ALSPAC data set but not in the QIMR cohort. Four SNPs located within the *AR* gene and six SNPs in high linkage disequilibrium (LD) with these displayed little evidence of association with the 2D:4D measures (best $p = 0.04$ for rs4456006). Similarly, there was no strong evidence for association between SNPs in the aromatase gene (*CYP19A1* [MIM 107910]) and any of the 2D:4D measures (all $p > 0.01$). Additionally, there were no large signals present in the *HOXD* cluster of genes (best




Figure 1. Plot of the Locus Surrounding *LIN28B*

SNPs are plotted by position on chromosome 6 (build 36) against GWAS p values for average 2D:4D. SNP rs314277 is shown in dark red labeled with its p value in the discovery meta-analysis. Estimated recombination rates from HapMap are plotted in light blue to reflect local LD structure. The directly genotyped or imputed SNPs surrounding rs314277 are color coded to reflect their LD with rs314277 (according to pair-wise r^2 values from the HapMap CEU database). Genes and their direction of transcription are labeled at the bottom of the plot.

$p = 0.0015$ for rs2857533), nor in the *HOXA* cluster (all $p > .05$), which play important roles in limb development.

We also performed a chi-square test for heterogeneity to test whether the regression coefficients differed significantly between males and females. No variants exhibited significant differences in the magnitude of the regression coefficients between the sexes, although this may be

partially a consequence of the low power of these tests to detect interactions. A complete list of variants with p values less than 10^{-5} is displayed in Tables S4–S6.

We attempted to replicate the *LIN28B* association by genotyping both rs314277 and rs314276 in an additional 5129 children from the ALSPAC cohort (Table 2). Whereas rs314277 was strongly associated with 2D:4D in the

Table 2. Association Results for the SNP Markers rs314277 and rs314276 in the ALSPAC Discovery GWAS, the QIMR Discovery GWAS, the Combined Meta-Analysis, and the Replication Data

	MAF	Left 2D:4D			Right 2D:4D			Mean 2D:4D		
		Beta	SE	p Value	Beta	SE	p Value	Beta	SE	p Value
rs314277, Alleles A/C										
ALSPAC Discovery	0.14	0.54	0.17	1.90×10^{-3}	0.58	0.17	4.90×10^{-4}	0.56	0.16	3.17×10^{-4}
QIMR Discovery	0.14	0.74	0.19	1.50×10^{-4}	0.65	0.18	4.30×10^{-4}	0.70	0.17	2.40×10^{-5}
Combined		0.63	0.13	1.50×10^{-6}	0.61	0.12	8.20×10^{-7}	0.63	0.11	4.10×10^{-8}
Replication	0.15	0.42	0.01	2.75×10^{-5}	0.47	0.10	5.15×10^{-6}	0.44	0.09	1.53×10^{-6}
rs314276, Alleles A/C										
ALSPAC Discovery	0.34	0.37	0.13	4.05×10^{-3}	0.27	0.12	0.027	0.32	0.12	5.30×10^{-3}
QIMR Discovery	0.34	0.39	0.14	6.00×10^{-3}	0.33	0.13	0.014	0.36	0.12	3.00×10^{-3}
Combined		0.38	0.09	7.92×10^{-5}	0.30	0.09	1.01×10^{-3}	0.34	0.08	5.42×10^{-5}
Replication	0.34	0.21	0.07	5.34×10^{-3}	0.30	0.08	1.12×10^{-4}	0.26	0.07	2.26×10^{-4}

In each case, the minor allele is listed first in the table. Beta coefficients indicate the expected change in 2D:4D per additional minor allele.

replication cohort ($\beta = 0.44$; 95% confidence interval: 0.26–0.62; $p = 1.53 \times 10^{-6}$), rs314276 showed only nominal association ($p = 2.26 \times 10^{-4}$). Indeed, conditioning on rs314277 suggested that the association at rs314276 could be entirely explained by the signal at rs314277 ($p = 0.202$). The association between rs314277 and 2D:4D also remained after conditioning on height ($p = 1.64 \times 10^{-6}$). There was no evidence for interaction between genotype at rs314277 and sex in the replication cohort ($p = 0.31$).

Because 2D:4D has been hypothesized to reflect testosterone exposure in utero, we investigated a possible relationship between mother's genotype at rs314277 and offspring 2D:4D. Although there was some evidence for a relationship between maternal genotype and mean 2D:4D in offspring ($p = 0.05$), any hint of association disappeared after conditioning on the child's genotype ($p = 0.99$), suggesting that the mother's genotype at rs314277 did not directly influence the child's 2D:4D.

Finally, given the previous association between rs314277 and height, we explored whether any of the confirmed variants from 11 different GWAS of height and four GWAS of age at menarche were associated with 2D:4D from our meta-analysis, using the online GWAS catalog (see [Web Resources](#)). Apart from SNPs located at 6q21, the only other variants that showed nominal association with 2D:4D were rs4932217, near the gene *POLG* (MIM 174763; $p = 0.01$), and rs2292303, in the gene *NUP37* (MIM 609264; $p = 0.05$). There did not appear to be an overlap in the direction of effect when the loci from all these studies were compared (i.e., variants associated with increased height did not generally appear to be associated with variants that increased [or, conversely, decreased] 2D:4D). Similarly, SNPs in the 9q31 region that have previously been associated with age at menarche³¹ did not show association with mean 2D:4D either (all $p > 0.05$). A complete list of variants previously associated with height and age at menarche are presented in [Table S7](#), along with results from the current combined meta-analysis of mean 2D:4D.

Discussion

The ratio between an individual's second and fourth digit (2D:4D) is sexually dimorphic and has been frequently used as a noninvasive retrospective biomarker for prenatal androgen exposure.⁹ 2D:4D has been shown to correlate with a wide range of diseases and physiological and psychological traits, including autism,¹⁰ attention deficit disorder,¹¹ fertility,¹² myocardial infarction,¹³ visuo-spatial ability,¹⁴ homosexuality,^{15,16} athletic performance,¹⁷ and age at menarche.¹⁸ In this study, we identified a variant located within intron 2 of the *LIN28B* gene, rs314277, which was robustly associated with 2D:4D.

LIN28B is the human ortholog of a gene that regulates developmental timing in *Caenorhabditis elegans*. Its

product, LIN28-B, is an RNA-binding protein that interacts directly with let-7 precursors, preventing their processing to become mature miRNAs.³² Polymorphisms within *LIN28B* have previously been associated with height³³ and age at menarche in females.^{28,34,35} Most recently, Viswanathan et al. demonstrated that activation of LIN28B promotes neoplastic transformation and is associated with aggressive forms of human malignancy.³⁶ Although it is unclear at present how variation in *LIN28B* might influence 2D:4D, it is noteworthy that the gene is highly expressed in the testis, placenta, and fetal liver,^{34,37} suggesting that *LIN28B* might influence 2D:4D early in development.

The association between rs314277 in *LIN28B* and 2D:4D is particularly interesting because the minor allele has previously been associated with increased height³³ and delayed menarche in females.³⁵ The direction of effect in these studies is consistent with the overall correlation between the variables, because girls who experience menarche later tend to be taller as adults than girls who reach puberty earlier.³⁸ Similarly, a recent study of 2D:4D and age at menarche found that girls with low right-hand 2D:4D (although not left-hand 2D:4D) tended to experience delayed menarche.¹⁸ In the present study, however, the minor allele at rs314277 was associated with *increased* 2D:4D ratio, which is opposite to the direction predicted by these earlier reports. It is unclear why this might be the case, but it suggests that the relationship between 2D:4D, age at menarche, and height is complex. Given that the association between rs314277 and 2D:4D remained after conditioning on height, and given that no other SNPs that displayed association with height or with age at menarche showed convincing evidence of association with 2D:4D, it is unlikely that the effect of rs314277 is mediated through these other variables. It is interesting to note that LIN28B has two isoforms distinguished by the presence or truncation of a conserved cold-shock domain, which influences protein function.³⁸ It is possible that different isoforms of LIN28B and hence different biological pathways might influence 2D:4D, height, and age at menarche, helping to explain the interesting pattern of correlations.

Our study is not the only report to have questioned the specificity of 2D:4D as a proxy for androgen exposure. In fact, the relationship between 2D:4D and prenatal androgen levels has been directly examined in only one small sample ($n = 33$), and no significant relationship between testosterone levels within the amniotic fluid (at ~17 wks gestation) and 2D:4D (at 24 mo) was found, but a correlation was found between 2D:4D and the ratio of fetal testosterone to estradiol,^{4,39} suggesting that the relationship between 2D:4D and circulating sex steroids might be more complex than previously hypothesized. In this study, we did not find evidence of a relationship between genetic variants in or near the androgen receptor and 2D:4D, although it is unclear how well the SNPs in this study might tag CAG repeat expansion at the androgen receptor.

In conclusion, we have demonstrated that a variant within the *LIN28B* gene is associated with 2D:4D and is important in the early development of the hands. The same variant is associated with height and age at menarche, but the direction of the association with 2D:4D was in the opposite direction to that predicted from an earlier study of 2D:4D and age at menarche. Our result suggests that the relationship between 2D:4D, early development, and fetal androgen exposure is likely to be more complex than previously appreciated.

Supplemental Data

Supplemental Data include seven tables and five figures and can be found with this article online at <http://www.ajhg.org>.

Acknowledgments

We are extremely grateful to all of the families who took part in the ALSPAC study, the midwives for their help in recruiting them, and the whole ALSPAC team. We thank John Manning for advice with the set up of the 2D:4D measurement, and we thank Jeremy Horwood and Andrea Waylen for help with implementation in the assessment clinics. The UK Medical Research Council, the Wellcome Trust, and the University of Bristol provide core support for ALSPAC and the MRC CAiTE (MRC 90600705). We thank the Sample Logistics and Genotyping facilities at the Wellcome Trust Sanger Institute for generating the ALSPAC genome-wide association data. D.M.E. and this work were supported by a Medical Research Council New Investigator Award (MRC G0800582 to D.M.E.).

We thank the twins and their families from the QIMR Brisbane Adolescent Twin Study for their participation; Dixie Statham, Ann Eldridge, Marlene Grace, and Kerrie McAloney for sample collection; Lisa Bowdler, Sara Smith, and Steven Crooks for DNA processing; and David Smyth, Harry Beeby, and Daniel Park for IT support. We thank the Australian National Health and Medical Research Council (NHMRC; grants 241944, 339462, 389927, 389875, 389891, 389892, 389938, 443036, 442915, 442981, 496739, 552485, and 552498) and the Australian Research Council (A7960034, A79906588, A79801419, DP0212016, and DP0343921) for funding the QIMR Brisbane Adolescent Twin Study. Statistical analyses were carried out on the Genetic Cluster Computer, which is financially supported by the Netherlands Scientific Organization (NWO 480-05-003). S.E.M., D.R.N., and G.W.M. are supported by the NHMRC Fellowship Scheme.

Received: November 26, 2009

Revised: January 27, 2010

Accepted: February 12, 2010

Published online: March 18, 2010

Web Resources

The URLs for data presented herein are as follows:

Genetic Cluster Computer, <http://www.geneticcluster.org>

KBIOSCIENCES, <http://www.kbioscience.co.uk>

MACH and MACH2QTL, <http://www.sph.umich.edu/csg/abecasis/MACH/download>

METAL, <http://www.sph.umich.edu/csg/abecasis/metal/>

Online GWAS Catalogue (accessed January 15, 2010), <http://www.genome.gov/gwastudies/>

References

1. Phelps, V.R. (1952). Relative index finger length as a sex-influenced trait in man. *Am. J. Hum. Genet.* 4, 72–89.
2. Ecker, A. (1875). Einige Bemerkungen über einen schwankenden charakter in der hand des menschen. *Archiv für Anthropologie* 8, 67–74.
3. Wilson, G.D. (1983). Finger length as an index of assertiveness in women. *Pers. Individ. Dif.* 4, 111–112.
4. Manning, J.T. (2002). Digit Ratio: A pointer to fertility, behavior, and health (New Brunswick: Rutgers University Press).
5. Manning, J.T., Scutt, D., Wilson, J., and Lewis-Jones, D.I. (1998). The ratio of 2nd to 4th digit length: a predictor of sperm numbers and concentrations of testosterone, luteinizing hormone and oestrogen. *Hum. Reprod.* 13, 3000–3004.
6. Malas, M.A., Dogan, S., Evcil, E.H., and Desdicioglu, K. (2006). Fetal development of the hand, digits and digit ratio (2D:4D). *Early Hum. Dev.* 82, 469–475.
7. McIntyre, M.H., Ellison, P.T., Lieberman, D.E., Demerath, E., and Towne, B. (2005). The development of sex differences in digital formula from infancy in the Fels Longitudinal Study. *Proc. Biol. Sci.* 272, 1473–1479.
8. McIntyre, M.H. (2006). The use of digit ratios as markers for perinatal androgen action. *Reprod. Biol. Endocrinol.* 4.
9. Voracek, M., and Loibl, L.M. (2009). Scientometric analysis and bibliography of digit ratio (2D:4D) research, 1998–2008. *Psychol. Rep.* 104, 922–956.
10. Manning, J.T., Baron-Cohen, S., Wheelwright, S., and Sanders, G. (2001). The 2nd to 4th digit ratio and autism. *Dev. Med. Child Neurol.* 43, 160–164.
11. Stevenson, J.C., Everson, P.M., Williams, D.C., Hipskind, G., Grimes, M., and Mahoney, E.R. (2007). Attention deficit/hyperactivity disorder (ADHD) symptoms and digit ratios in a college sample. *Am. J. Hum. Biol.* 19, 41–50.
12. Bang, A.K., Carlsen, E., Holm, M., Petersen, J.H., Skakkebaek, N.E., and Jørgensen, N. (2005). A study of finger lengths, semen quality and sex hormones in 360 young men from the general Danish population. *Hum. Reprod.* 20, 3109–3113.
13. Manning, J.T., and Bundred, P.E. (2000). The ratio of 2nd to 4th digit length: a new predictor of disease predisposition? *Med. Hypotheses* 54, 855–857.
14. Collaer, M.L., Reimers, S., and Manning, J.T. (2007). Visuospatial performance on an internet line judgment task and potential hormonal markers: sex, sexual orientation, and 2D:4D. *Arch. Sex. Behav.* 36, 177–192.
15. Manning, J.T., and Robinson, S.J. (2003). 2nd to 4th digit ratio and a universal mean for prenatal testosterone in homosexual men. *Med. Hypotheses* 61, 303–306.
16. Rahman, Q., and Wilson, G.D. (2003). Sexual orientation and the 2nd to 4th finger length ratio: evidence for organising effects of sex hormones or developmental instability? *Psychoneuroendocrinology* 28, 288–303.
17. Hönekopp, J., T Manning, J., and Müller, C. (2006). Digit ratio (2D:4D) and physical fitness in males and females: Evidence for effects of prenatal androgens on sexually selected traits. *Horm. Behav.* 49, 545–549.

18. Matchock, R.L. (2008). Low digit ratio (2D:4D) is associated with delayed menarche. *Am. J. Hum. Biol.* *20*, 487–489.
19. Gobrogge, K.L., Breedlove, S.M., and Klump, K.L. (2008). Genetic and environmental influences on 2D:4D finger length ratios: a study of monozygotic and dizygotic male and female twins. *Arch. Sex. Behav.* *37*, 112–118.
20. Medland, S.E., and Loehlin, J.C. (2008). Multivariate genetic analyses of the 2D:4D ratio: examining the effects of hand and measurement technique in data from 757 twin families. *Twin Res. Hum. Genet.* *11*, 335–341.
21. Paul, S.N., Kato, B.S., Cherkas, L.F., Andrew, T., and Spector, T.D. (2006). Heritability of the second to fourth digit ratio (2d:4d): A twin study. *Twin Res. Hum. Genet.* *9*, 215–219.
22. Voracek, M., and Dressler, S.G. (2007). Digit ratio (2D:4D) in twins: heritability estimates and evidence for a masculinized trait expression in women from opposite-sex pairs. *Psychol. Rep.* *100*, 115–126.
23. Manning, J.T., Bundred, P.E., Newton, D.J., and Flanagan, B.F. (2003). The 2nd to 4th digit ratio and variation in the androgen receptor gene. *Evol. Hum. Behav.* *24*, 399–405.
24. Golding, J., Pembrey, M., Jones, R., and ALSPAC Study Team. (2001). ALSPAC—the Avon Longitudinal Study of Parents and Children. I. Study methodology. *Paediatr. Perinat. Epidemiol.* *15*, 74–87.
25. Wright, M.J., and Martin, N. (2004). Brisbane Adolescent Twin Study: outline of study methods and research projects. *Aust. J. Psychol.* *56*, 65–78.
26. Timpson, N.J., Tobias, J.H., Richards, J.B., Soranzo, N., Duncan, E.L., Sims, A.M., Whittaker, P., Kumanduri, V., Zhai, G., Glaser, B., et al. (2009). Common variants in the region around Osterix are associated with bone mineral density and growth in childhood. *Hum. Mol. Genet.* *18*, 1510–1517.
27. Medland, S., Nyholt, D., Painter, J., McEvoy, B., McRae, A., Zhu, G., Gordon, S., Wray, N., Ferreira, M., Wright, M., et al. Common variants in the Trichohyalin gene are associated with straight hair in Europeans. *Am. J. Hum. Genet.* *85*, 750–755.
28. Ong, K.K., Elks, C.E., Li, S., Zhao, J.H., Luan, J., Andersen, L.B., Bingham, S.A., Brage, S., Smith, G.D., Ekelund, U., et al. (2009). Genetic variation in LIN28B is associated with the timing of puberty. *Nat. Genet.* *41*, 729–733.
29. Abecasis, G.R., Cherny, S.S., Cookson, W.O., and Cardon, L.R. (2002). Merlin—rapid analysis of dense genetic maps using sparse gene flow trees. *Nat. Genet.* *30*, 97–101.
30. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., Maller, J., Sklar, P., de Bakker, P.I., Daly, M.J., and Sham, P.C. (2007). PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* *81*, 559–575.
31. Perry, J.R., Stolk, L., Franceschini, N., Lunetta, K.L., Zhai, G., McArdle, P.F., Smith, A.V., Aspelund, T., Bandinelli, S., Boerwinkle, E., et al. (2009). Meta-analysis of genome-wide association data identifies two loci influencing age at menarche. *Nat. Genet.* *41*, 648–650.
32. Viswanathan, S.R., Daley, G.Q., and Gregory, R.I. (2008). Selective blockade of microRNA processing by Lin28. *Science* *320*, 97–100.
33. Lettre, G., Jackson, A.U., Gieger, C., Schumacher, F.R., Berndt, S.I., Sanna, S., Eyheramendy, S., Voight, B.F., Butler, J.L., Guiducci, C., et al. Diabetes Genetics Initiative, FUSION, KORA, Prostate, Lung Colorectal and Ovarian Cancer Screening Trial, Nurses' Health Study, SardiNIA. (2008). Identification of ten loci associated with height highlights new biological pathways in human growth. *Nat. Genet.* *40*, 584–591.
34. Sulem, P., Gudbjartsson, D.F., Rafnar, T., Holm, H., Olafsdottir, E.J., Olafsdottir, G.H., Jonsson, T., Alexandersen, P., Feenstra, B., Boyd, H.A., et al. (2009). Genome-wide association study identifies sequence variants on 6q21 associated with age at menarche. *Nat. Genet.* *41*, 734–738.
35. He, C., Kraft, P., Chen, C., Buring, J.E., Pare, G., Hankinson, S.E., Chanock, S.J., Ridker, P.M., Hunter, D.J., and Chasman, D.I. (2009). Genome-wide association studies identify loci associated with age at menarche and age at natural menopause. *Nat. Genet.* *41*, 724–728.
36. Viswanathan, S.R., Powers, J.T., Einhorn, W., Hoshida, Y., Ng, T.L., Toffanin, S., O'Sullivan, M., Lu, J., Phillips, L.A., Lockhart, V.L., et al. (2009). Lin28 promotes transformation and is associated with advanced human malignancies. *Nat. Genet.* *41*, 843–848.
37. Guo, Y., Chen, Y., Ito, H., Watanabe, A., Ge, X., Kodama, T., and Aburatani, H. (2006). Identification and characterization of lin-28 homolog B (LIN28B) in human hepatocellular carcinoma. *Gene* *384*, 51–61.
38. Onland-Moret, N.C., Peeters, P.H., van Gils, C.H., Clavel-Chapelon, F., Key, T., Tjønneland, A., Trichopoulou, A., Kaaks, R., Manjer, J., Panico, S., et al. (2005). Age at menarche in relation to adult height: the EPIC study. *Am. J. Epidemiol.* *162*, 623–632.
39. Lutchmaya, S., Baron-Cohen, S., Raggatt, P., Knickmeyer, R., and Manning, J.T. (2004). 2nd to 4th digit ratios, fetal testosterone and estradiol. *Early Hum. Dev.* *77*, 23–28.